

Assessing the burden of rheumatic heart disease among refugee children: a call to action

Gabriele Rossi¹, Vivian Sau Wai Lee²

¹ Médecins Sans Frontières, Operational Center, Brussels, Belgium

² Médecins Sans Frontières Hong Kong, Sai Wan, Hong Kong

In a recent editorial in *The Lancet*, Langlois and colleagues are adamant in advocating for a better and more universal access to healthcare for refugees landing in Europe [1]. The arguments of equity, social justice and cost-effectiveness are clearly explained in their comment.

Among the different suggestions to improve the refugees' health, the authors invite the policy makers to opt for strategies favoring the "provision of preventive care, including primary and secondary prevention of cardiovascular disease" which "could generate savings for health-care systems by alleviating the burden of stroke, and myocardial infarction".

Their advice, geared towards the betterment of the refugees' adult population, needs to be further complemented by a strong call for providing primary and secondary preventive care for acquired cardiac diseases in refugee children.

While acute rheumatic fever (ARF) has essentially vanished from industrialised countries since the latter half of the 20th century, the condition and its major sequel, rheumatic heart disease (RHD) remains an important public health concern in poor, developing and war-torn countries.

Poverty, inadequate and disrupted primary healthcare systems are major contributors to the persistence and resurgence of ARF/RHD in these countries [2].

The estimate of the global burden of RHD has been recently established at 33 million existing cases, which is higher than the previous conservative 15.6 to 19.6 million cases in 2005 [3]. Emerging echocardiographic data following the seminal work of Marijon and colleagues [4], have contributed in revising these data, suggesting for a higher prevalence of subclinical RHD. In particular, according to a recent meta-analysis by Rothenbuler et al [5], among

The current enormous refugee crisis in Europe represents a paradigmatic translational shift: the health systems of the wealthy countries are now called to deal, in a substantial way, with the poor people and their diseases.

children aged below 18 years from 37 pooled countries, the prevalence of clinically silent rheumatic heart disease (21.1 per 1000 people, 95% confidence interval CI 14.1–31.4) would be about seven to eight times higher than that of clinically manifest disease (2.7 per 1000 people, 95% CI 1.6–4.4). South Asia and some African countries have very high prevalence data [5].

Although the significance of subclinical RHD is still not clear, follow up studies indicate that most Subclinical Definite RHD (according to World Heart Federation Criteria) [6] and at least a proportion of Borderline RHD, are likely to be true RHD requiring secondary prophylaxis [7].

To tackle the evolution of RHD, specific control programs based on the concurrent development of disease registers along with the consistent delivering of benzathine penicillin G injections, need to be strengthened by a decentralized echocardi-based active case finding activities, comprehensively integrated into the existing primary health care services.

In this sense, the World Heart Federation's non-communicable disease action plan is calling for a 25% reduction in premature mortality from RHD by the year 2025 ("25 by 25") [6].





Photo: Médecins Sans Frontières assists Syrian refugees in Roske, at the Hungary-Serbia border. Among the refugees arriving in Europe, half of them are children. Courtesy of Ana Lemos/MSF (2015).

Unfortunately, this cost-effective strategy has not been routinely implemented in low-income and middle-income countries, because of the structural weaknesses of their health systems.

The current enormous refugee crisis in Europe represents a paradigmatic translational shift: the health systems of the wealthy countries are now called to deal, in a substantial way, with the poor people and their diseases.


The Syrian children, suffering from more than five years of displacement, have been forced to live in very dire conditions. Being exposed to the effects of the detrimental combination of seasonal harsh cold weather, overcrowding, poverty and very limited access to healthcare [8], they hold the classic risky profile for the development of ARF and subsequent RHD [9].

With an expected echocardiography-based screening detection rate of around 2%, many migrant children landing in Europe (Syrian, Afghan, Eritrean, Bangladeshi, and many others) could be timely identified and placed under prophylactic treatment for RHD. In 2015, more than 1 million asylum seekers arrived in Europe and half of them were children: the figure of 10 000 diagnosed subclinical and clinical RHD could be anticipated. Many more refugees are expected to land in Europe in 2016.

Pilot studies aimed at proving the validity of these assumptions, while assessing the prevalence of RHD among refugee children, are urgently warranted.

Decentralised diagnostic services for ARF/RHD, framed as routine screening through point-of care technologies (echocardiography and antigen tests for the rapid diagnosis of group A streptococcal pharyngitis) should be then considered at migrant/refugee pediatric community level.

In view of the fact that severe RHD is lethal in the absence of surgical treatment, with a death toll of 275 000/year [10], the same arguments of social justice, equity and cost-effectiveness mentioned by Langlois and colleagues should apply here, especially for the silent and marginalized population of refugee and migrant children, often arriving in Europe unaccompanied and defenseless. Europe has the infrastructures, human and financial resources, and knowledge to act accordingly. It must not turn a blind eye to them.



Acknowledgements: We would like to thank the Médecins Sans Frontières (MSF) Associative Boards of Italy and Hong Kong, as well as the MSF Operational Center Brussels (MSF OCB), for fostering and supporting a continued and meaningful medical action in favor of the refugee populations across Europe and South-East Asia.

Funding: None.

Disclaimer: None.

Authorship declaration: GR contributed with the idea and the content of the manuscript. VSW contributed with the editing of the manuscript.

Competing interests: The authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare no conflict of interest.

REFERENCES

- 1 Langlois EV, Haines A, Tomson G, Ghaffar A. Refugees: towards better access to healthcare services. *Lancet*. 2016;387:319-21. [Medline:26842434doi:10.1016/S0140-6736\(16\)00101-X](https://doi.org/10.1016/S0140-6736(16)00101-X)
- 2 Omurzakova NA, Yamano Y, Saatova GM, Mirzakhanova MI, Shukurova SM, Kydyralieva RB, et al. High incidence of rheumatic fever and rheumatic heart disease in the republics of Central Asia. *Int J Rheum Dis*. 2009;12:79-83. [Medline:20374323doi:10.1111/j.1756-185X.2009.01388.x](https://doi.org/10.1111/j.1756-185X.2009.01388.x)
- 3 Global Burden of Disease Study 2013 Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2015;386:743-800. [Medline:26063472doi:10.1016/S0140-6736\(15\)60692-4](https://doi.org/10.1016/S0140-6736(15)60692-4)

- 4 Marijon E, Ou P, Celermajer DS, Ferreira B, Mocumbi AO, Jani D, et al. Prevalence of rheumatic heart disease detected by echocardiographic screening. *N Engl J Med.* 2007;357:470-6. [Medline:17671255](#)[doi:10.1056/NEJMoa065085](#)
- 5 Rothenbühler M, O'Sullivan CJ, Stortecky S, Stefanini GG, Spitzer E, Estill J, et al. Active surveillance for rheumatic heart disease in endemic regions: a systematic review and meta-analysis of prevalence among children and adolescents. *Lancet Glob Health.* 2014;2:e717-26. [Medline:25433627](#)[doi:10.1016/S2214-109X\(14\)70310-9](#)
- 6 World Heart Federation. World Heart Federation RF and RHD goals: 25x25<25. Available: <http://www.world-heart-federation.org/what-we-do/rheumatic-heart-diseasenetwork/> world-heart-federation-rf-and-rhd-goals-25x25. Accessed: 21 March 2016.
- 7 Zühlke L, Engel ME, Lemmer CE, van de Wall M, Nkepu S, Meiring A, et al. The natural history of latent rheumatic heart disease in a 5 year follow-up study: a prospective observational study. *BMC Cardiovasc Disord.* 2016;16:46. [Medline:26892774](#)[doi:10.1186/s12872-016-0225-3](#)
- 8 UN News Center. Syria: UNICEF plans to reach 2.6 million children with winter supplies and cash assistance. Available: <http://www.un.org/apps/news/story.asp?NewsID=52757#.VrGmAjYcYG7J>. Accessed: 22 March 2016.
- 9 Islam AK, Majumder AA. Rheumatic fever and rheumatic heart disease in Bangladesh: a review. *Indian Heart J.* 2016;68:88-98. [Medline:26896274](#)[doi:10.1016/j.ihj.2015.07.039](#)
- 10 GBD 2013 Mortality and Causes of Death Collaborators. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet.* 2015;385:117-71. [Medline:25530442](#)[doi:10.1016/S0140-6736\(14\)61682-2](#)

Correspondence to:

Gabriele Rossi Rossi, MD, MPH
Via Como 17
20095 Cusano Milanino
Italy
gab.rossi@tiscali.it