

Nutrition: the missing link in the battle against microbial resistance?

Stefan A Unger^{1,2*}, Henry Mark^{3*}, Claudia Pagliari^{4*}

¹ University of Edinburgh, Department of Child Life and Health, Edinburgh, UK

² Royal Hospital for Sick Children, Department of Respiratory Medicine, Edinburgh, UK

³ International Development Consultant in Nutrition, Nottingham, UK

⁴ University of Edinburgh, Usher Institute of Population Health Sciences and Informatics, Edinburgh, UK

* Equal authorship

Active and preventative patient-centred nutritional interventions within primary health care and communities have a key role to play in addressing the global challenge of AMR and should be considered an essential element of strategies to provide antibiotic stewardship and security.

The growing threat posed by antimicrobial resistance (AMR) continues to attract scientific and policy attention [1,2]. A number of recent publications have reignited the debate on striking a balance between the use of antibiotics as a powerful public health tool [3] and the significant risks posed by AMR, including an increase in mortality from drug-resistant strains of common bacterial infections in children with HIV, TB and malaria [4].

Rochford and others recently called for an international inter-agency effort to strengthen global governance of AMR [1].

They suggest setting up a multi-stakeholder Global Steering Board, within existing organisational structures, to develop a legal agreement on international standards and norms with the following four key objectives: 1) ensuring the appropriate use of antibiotics in humans and animals, 2) eradicating untreated effluent, 3) strengthening infection prevention and control measures such as water, sanitation, and hygiene programmes; and 4) providing appropriate access to a regulated supply of quality-assured affordable antimicrobials. We agree with all of these points, but would argue that an important contribution to AMR prevention should be added to these objectives: the promotion of good nutrition. Here we draw on indirect evidence and theory to support this proposition.

Following previous observations in Ethiopia of a possible reduction in childhood mortality resulting from antibiotic interventions tackling trachoma, the New England Journal of Medicine published details of a multi-country cluster-randomised trial in Sub-Saharan Africa, providing twice-yearly mass administration of azithromycin vs placebo to children aged 1 to 59 months [3]. A total of 190 238 children from 1533 randomized communities in Malawi, Niger and Tan-



Photo: iStock.


risks of processed food supplementation [11]. Even when targeting children presenting with an acute illness, short-term supplementation with small quantities of micronutrient fortified lipid-based supplements and ready-to-use therapeutic foods appears to have only limited effects on subsequent growth, future severe malnutrition or illness episodes [11].

Despite these complexities, it is reasonable to argue that better nourished children will be less susceptible to infectious diseases and thus less likely to require antibiotics. A similar logic was given by Laxminarayan et al in their paper ‘Antibiotic resistance – the need for global solutions’ [12], who speculate that the effects of AMR are likely to be higher in vulnerable groups, including those who are malnourished. *We therefore propose that strategies for global antibiotic stewardship should include interventions to both tackle malnutrition and promote optimal nutrition as an upstream influence on both infectious disease and AMR.*

Given the need for AMR strategies to integrate multiple disease prevention tactics, including interventions to improve hygiene and child feeding practices, holistic, person-centred, multi-layered approaches are most likely to deliver lasting results. We suggest that all health care encounters, at community, primary or secondary levels provide a unique opportunity to: 1) offer nutritional education, to tackle low awareness or cultural barriers [13]; 2) provide targeted nutritional interventions, including treatment for malnourished children [13], 3) provide guidance on hygiene promotion [14] and symptom recognition for severe infectious diseases [15]; and 4) diagnose and treat infections appropriately [16]. Such a combined approach may provide mothers and carers with the tools required to reduce the incidence of infectious disease, decrease time to recovery and lower the risk of re-infection, thus reducing the need for antibiotics, and, by extension, helping to combat AMR.

Mobile phone-based (mHealth) tools have potential to support the delivery of such holistic interventions as well as to collect data on nutritional and infection indicators to guide the targeting of interventions and enable smarter AMR management [17]. While holding much promise for improving population health, such multifaceted approaches require strong country-level commitment and community engagement [18]. At the same time, global political solutions and agreements are needed to ensure the fair and secure distribution of food, rather than focusing on crisis-driven supplementation.

We strongly recommend that nutrition be considered as an essential element of strategies to provide antibiotic stewardship and security. Active and preventative, patient-centred nutritional interventions within primary health care and communities have a key role to play in addressing the global challenge of AMR, supported by further research to unpick the complex relationship between nutrition, infection and AMR.



Acknowledgements: The authors would like to thank Catriona-Robyn MacDonald for her invaluable help with the design of the figure.

Funding: None.

Authorship contributions: All authors contributed equally to the development of the idea and the design and preparation of the manuscript.

Competing interests: The authors completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available upon request from the corresponding author), and declare no conflicts of interest.

REFERENCES

- 1 Rochford C, Sridhar D, Woods N, Saleh Z, Hartenstein L, Ahlawat H, et al. Global governance of antimicrobial resistance. *Lancet*. 2018;391:1976-8. [Medline:29864011](#) [doi:10.1016/S0140-6736\(18\)31117-6](#)
- 2 Zetts RM, Stoesz A, Smith BA, Hyun DY. Outpatient Antibiotic use and the need for Increased antibiotic stewardship efforts. *Pediatrics*. 2018;141:e20174124. [Medline:29793986](#) [doi:10.1542/peds.2017-4124](#)
- 3 Keenan JD, Bailey RL, West SK, Arzika AM, Hart J, Weaver J, et al. Azithromycin to reduce childhood mortality in sub-Saharan Africa. *N Engl J Med*. 2018;378:1583-92. [Medline:29694816](#) [doi:10.1056/NEJMoa1715474](#)
- 4 Manning ML, Septimus EJ, Ashley ESD, Cosgrove SE, Fakhri MG, Schweon SJ, et al. Antimicrobial stewardship and infection prevention – leveraging the synergy: A position paper update. *Am J Infect Control*. 2018;46:364-8. [Medline:29592832](#) [doi:10.1016/j.ajic.2018.01.001](#)
- 5 The Lancet Infectious Diseases. Azithromycin to prevent deaths in children. *Lancet Infect Dis*. 2018;18:583. [Medline:29856346](#) [doi:10.1016/S1473-3099\(18\)30300-1](#)
- 6 Elvidge CD, Sutton PC, Ghosh T, Tuttle BT, Baugh KE, Bhaduri B, et al. A global poverty map derived from satellite data. *Comput Geosci*. 2009;35:1652-60. [doi:10.1016/j.cageo.2009.01.009](#)

- 7 UNICEF Expanded Global Databases_Wasting 2018. Available: <https://data.unicef.org/data/malnutrition/>. Accessed: 3rd July 2018.
- 8 Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet*. 2013;382:427-51. [Medline:23746772](#) [doi:10.1016/S0140-6736\(13\)60937-X](#)
- 9 Mark H, Been J, Sonko B, Faal A, Ngum M, Hasan J, et al. Nutritional status and disease severity in children acutely presenting to a primary health clinic in rural Gambia. *BMC Public Health*. 2019.14:e1002377.
- 10 Schaible UE, Kaufmann SH. Malnutrition and infection: complex mechanisms and global impacts. *PLoS Med*. 2007;4:e115. [Medline:17472433](#) [doi:10.1371/journal.pmed.0040115](#)
- 11 Unger SA, Drammeh S, Hasan J, Ceesay K, Sinjanka E, Beyai S, et al. Impact of fortified versus unfortified lipid-based supplements on morbidity and nutritional status: A randomised double-blind placebo-controlled trial in ill Gambian children. *PLoS Med*. 2017;14:e1002377. [Medline:28809926](#) [doi:10.1371/journal.pmed.1002377](#)
- 12 Laxminarayan R, Duse A, Wattal C, Zaidi AK, Wertheim HF, Sumpradit N, et al. Antibiotic resistance – the need for global solutions. *Lancet Infect Dis*. 2013;13:1057-98. [Medline:24252483](#) [doi:10.1016/S1473-3099\(13\)70318-9](#)
- 13 Fernandez-Barres S, Garcia-Barco M, Basora J, Martinez T, Pedret R, Arija V. The efficacy of a nutrition education intervention to prevent risk of malnutrition for dependent elderly patients receiving Home Care: A randomized controlled trial. *Int J Nurs Stud*. 2017;70:131-41. [Medline:28273591](#) [doi:10.1016/j.ijnurstu.2017.02.020](#)
- 14 Mills J, Cumming O. The impact of water, sanitation and hygiene on key health and social outcomes: Review of evidence. Available: https://www.unicef.org/wash/files/The_Impact_of_WASH_on_Key_Social_and_Health_Outcomes_Review_of_Evidence.pdf. Accessed: 1st January 2019.
- 15 Geldsetzer P, Williams TC, Kirolos A, Mitchell S, Ratcliffe LA, Kohli-Lynch MK, et al. The recognition of and care seeking behaviour for childhood illness in developing countries: a systematic review. *PLoS One*. 2014;9:e93427. [Medline:24718483](#) [doi:10.1371/journal.pone.0093427](#)
- 16 Kallander K, Burgess DH, Qazi SA. Early identification and treatment of pneumonia: a call to action. *Lancet Glob Health*. 2016;4:e12-3. [Medline:26577842](#) [doi:10.1016/S2214-109X\(15\)00272-7](#)
- 17 Hossain M, Choudhury N, Adib Binte Abdullah K, Mondal P, Jackson AA, Walson J, et al. Evidence-based approaches to childhood stunting in low and middle income countries: a systematic review. *Arch Dis Child*. 2017;102:903. [Medline:28468870](#) [doi:10.1136/archdischild-2016-311050](#)
- 18 Lee SH, Nurmatov UB, Nwaru BI, Mukherjee M, Grant L, Pagliari C. Effectiveness of mHealth interventions for maternal, newborn and child health in low- and middle-income countries: Systematic review and meta-analysis. *J Glob Health*. 2016;6:010401. [Medline:26649177](#) [doi:10.7189/jogh.06.010401](#)

Correspondence to:

Dr Stefan A Unger
 Department of Child Life and Health
 University of Edinburgh
 20 Sylvan Place
 Edinburgh, EH9 1UW
 UK
sunger@staffmail.ed.ac.uk